

Symmetry and planar chirality measured with a log-polar transformation in a transmission electron microscope

A. H. Tavabi,¹ P. Rosi,² R. B. G. Ravelli,³ A. Gijsbers,³ E. Rotunno,² T. Guner,⁴ Y. Zhang,³ A. Roncaglia,⁵ L. Belsito,⁵ G. Pozzi,^{1,6} T. Denneulin,¹ G.C. Gazzadi,² M. Ghosh,⁷ R. Nijland,⁷ S. Frabboni,^{2,6} P. J. Peters,³ E. Karimi,⁴ P. Tiemeijer,⁷ R. E. Dunin-Borkowski,¹ and V. Grillo²

¹ Ernst Ruska-Centre for Microscopy and Spectroscopy with Electrons and Peter Grünberg Institute, Forschungszentrum Jülich, 52425 Jülich, Germany

² Centro S3, CNR - Istituto di Nanoscienze, 41125 Modena, Italy

³ Maastricht Multimodal Molecular Imaging Institute, Division of Nanoscopy, Maastricht University, Universiteitsingel 50, Maastricht 6229 ER, The Netherlands

⁴ Department of Physics, University of Ottawa, Ottawa, Ontario K1N 6N5, Canada

⁵ CNR - Istituto per la Microelettronica e i Microsistemi-CNR, 40129 Bologna, Italy

⁶ Dipartimento FIM, Università di Modena e Reggio Emilia, 41125 Modena, Italy

⁷ Thermo Fisher Scientific, PO Box 80066, 5600 KA Eindhoven, the Netherlands




Fig. S1 Experimental defocus series for a logarithmic spiral. The apparent chirality changes with defocus.